



Enhanced Model for Monitoring and Quality Control Using the Control Chart for Binomial Regression with Application

Mohamed S. Hamed ^{a,b*}, Shaimaa Y. Mohammed ^c,
Salah m. Mohamed ^d and Amal Mohamed Abdel Fattah ^d

^a Department of Business Administration, Gulf Colleges, KSA, Saudi Arabia.

^b Department of Statistics & Mathematics and Insurance, Faculties of Commerce, Benha University, Egypt.

^c Bader Institutes for Information's technology, Badr City, Egypt.

^d Department of Applied Statistics, Faculty of Graduate Studies for Statistical Research, Cairo University, Egypt.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajpas/2024/v26i11667>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124816>

Received: 05/08/2024

Accepted: 08/10/2024

Published: 19/10/2024

Original Research Article.

Abstract

The aim of this study is to use a new approach called the Shewhart control chart with binomial regression, but under this condition, there was a multicollinearity problem in binomial regression model, and solving this problem by principal component, binomial regression with control chart is a statistical method used in quality control to monitor and improve the quality of products or services. This method involves using binomial regression to model the relationship between an input variable and output variable, and then integrating

*Corresponding author: Email: mssh@gulf.edu.sa;

Cite as: Hamed, Mohamed S., Shaimaa Y. Mohammed, Salah m. Mohamed, and Amal Mohamed Abdel Fattah. 2024. "Enhanced Model for Monitoring and Quality Control Using the Control Chart for Binomial Regression With Application". *Asian Journal of Probability and Statistics* 26 (11):1-8. <https://doi.org/10.9734/ajpas/2024/v26i11667>.

control charts to detect any changes in the process over time. The study used another control chart with a new approach and use another mythology to treat multicollinearity problems, also we used missing values and draw control charts.

Keywords: Quality control; model monitoring; shewhart control chart; binomial regression model; Generalized Linear Model (GLM).

1 Introduction

Model monitoring and quality control are vital in statistical analysis and predictive modelling, as they are responsible for guaranteeing the dependability, precision, and resilience of the developed models. These procedures encompass ongoing assessment, validation, and enhancement of the models to uphold their efficacy in making forecasts and informed choices. However, (Bilen et al., 2007) have developed a comprehensive framework for controlling multivariate auto-correlated processes that rely on principal component analysis and multiple regression control charts. The framework begins by identifying critical process variables using principal component analysis. Next, the autoregressive error correction model's white noise series is used to create a control chart that can be used for ongoing process monitoring. So, organizations must use quality control methods to guarantee the quality of their goods and services. The quality characteristic to be monitored in the second stage of two-stage processes whose distribution is binomial is the subject of this study's proposed control chart. The deviance residual, which is simply the generalized log-likelihood ratio statistic derived from the generalized linear model, forms the basis of the suggested control chart. And suggest applying a new link function in a generalized linear model framework to determine the relationship between the first- and second-stage quality features. Regarding the average run length requirement, the performance of the suggested control chart with the new link function is contrasted with that under the conventional logit link function. Furthermore, a comparison is made between the suggested control chart's performance and that of the chart created using the original residuals under the new link function, as well as the conventional np-chart used to track the binomial quality characteristic during the second stage (Amiria et al. 2016).

This study concentrated on the statistical method of creating a regression residual control chart using regression analysis to track students' academic progress in higher education institutions. While the variability was being tracked using the Moving Range Residual Control Chart, in e moving range chart, there were three spots out of control and one out of control in the individual control chart. As a result, the out-of-control locations were eliminated, and new, updated control boundaries were determined (Rashid et al. 2013). But in (Yassin and Mohamed, 2022) used one useful and popular method for creating associations between pairs of endogenous and exogenous variables regression modelling. They typically experience multicollinearity, though. This work employs residual control charts on count data (Poisson regression) following the application of the ridge technique to address multicollinearity issues. But (Filho et al., 2016) created residual control charts with Poisson regression. And used the principal component to solve the multicollinearity issue and then created a control chart. We used average run length as the statistics to assess the control chart. To improve quality, control charts are crucial statistical quality control instruments. (Nancy et al., 2023) explained the method and graphical procedure known as statistical quality control aided process monitoring and control. For the desired level of quality. Standard Shewhart control charts and their variations and control charts that are impacted by associated independent variables for the process shift are two distinct ways to look at control charts. In statistical process control, control charts are effective tools for monitoring processes. When examining a process based on an exponential family distributed response variable (such as binary outcomes) along with a single explanatory variable, the generalized linear model (GLM) provides better estimates.

The generalized linear mode (GLM) 1 is introduced by Nelder and Wedderburn, so the binomial regression model is a specific state of a generalized linear model. In other words, the statistical relationship between one or more independent variables and dependent variables can be verified. Furthermore, three elementary parts create the basis of a generalized linear model: a systematic component created from predictor variables that yield a linear predictor; a link function that connects the random and systematic parts; and a random component represented by the binary response variable, an average vector, and an exponential distribution (Soares et al. 2006).

This paper aims to discuss the binomial regression with multicollinearity problems that exists, solve this problem by PCA draw the residual-based Shewhart control chart, and use the ARL as a performance measure.

The remaining sections of the paper provide a concise overview of the Binomial Regression Model and only one type of residual, namely ordinary raw residual, in Section 2. Section 3 introduces our proposed methodology, which utilizes two approaches: (i) the application of the Principal Component Formula to address multicollinearity issues, and (ii) the use of a Residual Control Chart (Shewhart). Additionally, control charts for the binomial Model are presented. Section 4 consists of simulation studies describing the algorithmic approach and basic program for generating the models. We encourage further discussion on these topics. Lastly, Section 5 serves as the conclusion.

2 Materials and Methods

2.1 Materials

2.1.1 Binomial regression model

In binomial regression, the dependent variable represents the percentage of success in n independent essays, each with a probability of occurring p . According to (Pardo et al., 2007) the Regression model follows a binomial distribution with index n and parameter P , i.e., $n y^\circ \sim B(n, p)$. The density function of binomial distribution is given by:

$$f(y^\circ, p) = \binom{n}{ny^\circ} p^{ny^\circ} (1-p)^{n-ny^\circ} \text{ where } 0 < p, y^\circ < 1. \quad (1)$$

(Amiri et al., 2016) stated that the logit link function for binomial Regression is

$$\text{Logit}(p_i) = \log\left(\frac{p_i}{(1-p_i)}\right)$$

(Soares et al., 2006) stated that the statistical regression model for binomial Regression is:

$$y^\circ_i = \varphi_0 + \varphi_1 x_{i1} + \varphi_2 x_{i2} + \dots + \varphi_k x_{ik}. \quad (2)$$

Where $i = 1, \dots, I$, and independent variable $x_i = (x_{i0}, x_{i1}, \dots, x_{ik})$, $\varphi = (\varphi_0, \varphi_1, \dots, \varphi_k)^t$. Furthermore,

$$p_i = p(x_i^t \varphi) = \frac{\exp(\sum_{j=0}^k \varphi_j x_{ij})}{1 + \exp(\sum_{j=0}^k \varphi_j x_{ij})}, \text{ where } j = 1, \dots, k. \quad (3)$$

A multicollinearity problem is known as when we have a high correlation between independent variables (x). The least squares estimates are unbiased when multicollinearity exists, but because of their enormous variances, they could not be very close to the true value. Principal components regression lowers the standard errors by biasing the regression estimates to some extent.

We have used in this study an ordinary raw residual given from (Dunn and Smyth 2018). The ordinary raw residual is as follows:

$$\mathfrak{R}_o = y^\circ - \hat{\mu} \quad (4)$$

where the test of the null hypothesis is $H_0: p = p_0$ against the alternative hypothesis $H_a: p \neq p_0$ (Amiria et al. 2016).

2.2 Methods

(I) Principal Component Analysis:

The definition of the principal component analysis refers to (Filho and Sant'Anna 2016). Principal Component Analysis (PCA) is a statistical technique used for dimensionality reduction with the goal of reducing the

complexity of the data, making it easier to visualize, interpret, and process while retaining the most important patterns. The formula for the i^{th} Principal Component (PC) score referring to observation x is given by:

$$z_i = x * u_i \tag{5}$$

Where z_i presents the i^{th} PC score, x is the observation u_i is the eigenvector corresponding to the i^{th} PC.

(ii) Control Charts (CCs)

The control charts are an important and most widely used tool for statistical quality control. Furthermore, the statistical quality control approach is used to placement the limits of control chart and realization the modifications for product or process refinement. In application cases, a lot of scenarios where the control charts are used for monitoring the product or process (Haridy and Benneyan 2024, Mongkoltawat et al. 2024).

2.3 Shewhart Control Chart (SCC) for a binomial regression model

The Shewhart Control Chart is a useful graphic statistical tool. (Yassin and Mohamed 2022) used to the Shewhart based on residual control chart (RCC), furthermore (SCC) is used to detect process shifts or changes reverse to EWEMA control chart. The control chart limits (lower and upper) are given by:

$$CL_{\mathfrak{R}} = E(\mathfrak{R}_n) \pm v\sqrt{Var(\mathfrak{R}_n)} \cong \pm v. \tag{6}$$

Where $\mathfrak{R} \sim N(0,1)$, the constant v refers to the amplitude between control limits that depend on the false alarm probability τ . So, we have used the Shewhart Control Chart to show the performance of the Principal Component.

2.4 The average run length

Average Run Length (ARL) is a statistical measurement tool for measuring the number of observations required before control chart signals an out-of-control process. It is used to evaluate the effectiveness of control charts to detect process shifts or changes. The ARL is based on the chart design and the parameters used to establish the control chart limits, and a longer ARL means a chart is slower to signal a process shift. A shorter ARL indicates that a chart is faster at identifying process changes, but may result in more false alarms (Phanyaem 2024, Wilson et al. 2024).

According to (Yassin and Mohamed 2022) if the control chart is in control the (ARL) is equal to $ARL_0 = 1/\hat{\alpha}$, but if control chart is out of control the (ARL) is equal to $ARL_1 = 1/(1 - \hat{\beta})$, where $\hat{\alpha}$ is the probability of false alarm (type I error) and $\hat{\beta}$ is the probability of true alarm (type two).

3 Results and Discussion

3.1 Simulation studies

By using the R program version 4.3.3 we draw the Shewhart control chart for binomial regression under the condition multicollinearity problem wase exists, we will use it to solve our problem by principal component. And, using the deviance residual to draw a chart, we are also going to employ the Shewhart Control Chart to show the performance of the principal component in different phases and phase II.

3.2 The steps of generating data

1. Put $N = 200, p = n$, and generate z where z_{ij} following a standard normal distribution.
2. Put the degree of correlation $\delta = 0.95$ in equation generating independent variables x by equation $x_{ij} = (1 - \delta^2)^{\frac{1}{2}}z_{ij} + \delta z_{ij}$, where $i = 1, \dots, n, j = 1, \dots, m$.
3. Then choosing φ under condition $\sum_{j=1}^m \varphi = 1$ and taking $\varphi_0 = 1.5$ from (Filho et al., 2016)

4. Generating dependent variable y° of binomial regression model following $n y^\circ \sim B(n, p)$.
5. Finally, generating p_i , furthermore, we are using it in step 4. Where p is given by: $p_i = \frac{\exp(x_i\theta)}{1+\exp(x_i\theta)}$, $i = 1, 2, \dots, n$.

Table 1. The correlation matrix

Variable	x_1	x_2	x_3	x_4
x_1	1.0000000	0.9973866	0.9973168	0.9986863
x_2	0.9973866	1.0000000	0.9965585	0.9984274
x_3	0.9973168	0.9965585	1.0000000	0.9986005
x_4	0.9986863	0.9984274	0.9986005	1.0000000

Table 1 shows that there are very strong positive (direct) correlations between all variables.

Table 2. The estimated coefficient of model

Terms	Estimate of ϕ	SE Coef	VIF	Z-value	P-value
Constant	-0.9178	0.1635		-5.615	1.97e-08
x_1	-2.5532	3.2164	381.6762	-0.794	0.42732
x_2	-9.4354	3.4138	328.4978	-2.764	0.00571
x_3	-0.3039	3.2785	365.6465	-0.92614	0.92614
x_4	11.3533	5.2941	1096.3540	2.145	0.03199
AIC	242.9				
CI	78.58528				

Table 2 presents the summary results of the binomial regression model with VIF value exceeding 1. This indicates the presence of high degree of multicollinearity from the analysis. The CI value is obtained at 78.58528. Furthermore, x_2 and x_4 were found to be statistically significant while x_1 and x_3 are not significant at 5% level.




Fig. 1. Phase one with sample size $n = 4$ and number of sample $m=50$ for in control process

3.3 Analysis of simulation study

In this section, we clarify the proposed monitoring technique through a simulation study, so we check our data by condition index, variance inflation factor, and correlation matrix to see if this data has a multicollinearity problem. So, the subgroups of the simulation study are in control, and Table 2 shows the value of the estimated coefficient of the model. Furthermore, in Fig. 1 phase one is obtained from the principal component, and the

average run length (ARL_0) is equal to 7.94 and the value of (ARL_1) is equal to 1.1. However, in Fig. 2 is obtained from residuals, and (ARL_0) is equal to 7.86 and the value of (ARL_1) is equal to 1, and Fig. 3 is the phase two control chart, so (ARL_1) value is equal to 1.16, but the value of (ARL_0) is equal to 8.98. Where the sample size is $N = 200$ and p is the number of independent variables, so $p = 4$, then we divide N into 50 subgroups and subgroups, and the sample size is equal to 4. So, if we make a comparison by ARL measurement between Fig. 1, Fig. 2, and Fig. 3, we find that the (ARL_0) value of Fig. 3 is better than Figs. 1 and 2 because the (ARL_0) value of the ARL is the largest value, but in (ARL_1) Fig. 2 is the best value because it is the smallest value. So if we do a comparison between (Yassin and Mohamed 2022) and this study we found our methodology is easy to use and we use PC method instead of Yassin and Mohamed method to solve multicollinearity problem and monitor processes and procedures our data through Shewhart control chart and obtained on ARL because in Yassin and Mohamed doing three k control charts and every chart should be calculate ARL so they have 6 chart for one type put her we have two charts, so the best chart in this study for ARL_0 is Fig. 3 but in ARL_1 is Fig. 2.




Fig. 2. Phase one for binomial principal component residual based on control chart




Fig. 3. Phase two for binomial principal component residual based on control chart

4 Conclusions

All binomial control charts are in control. Furthermore, we present a new plan combining binomial regression and principal component analysis. Furthermore, the binomial principal component residual based on control chart (BPCR) is used to monitor data processes. A lot of research has used the Shewhart control chart, but Yassin and Mohamed used Shewhart based on a residual control chart with Poisson regression, so the current paper proposes a different strategy in terms of choosing a different model with a different treatment method. Furthermore, we generated z following standard normal distribution and use, $x_{ij} = (1 - \delta^2)^{\frac{1}{2}}z_{ij} + \rho z_{ij}$ with a degree of correlation equal to 0.95, then choosing φ finally generating dependent variable following binomial distribution then check the data by VIF, CI, and correlation matrix to show if this data have multicollinearity problem then do PCA, finally divided our data to subgroup and calculating ARL by. We found this suggested method is easy to use, doesn't take a lot of charts, and also can use different residuals to draw a control chart and calculate ARL if the control chart is in control the (ARL) is equal to $ARL_0 = 1/\hat{\alpha}$, but if control chart is out of control the (ARL) is equal to $ARL_1 = 1/(1 - \hat{\beta})$, however, it must be the largest value of ARL_0 because this value is the best value, but in ARL_1 should choose a small value. So if we make comparison, we found in Yassin and Mohamed the value is high for in control and out of control, but in our study, the value out of control is the best value because the value is smaller than Yassin and Mohamed, but the value of Yassin and Mohamed for in control is the best because it is greater than our study, that is why we say that our study gives good results compared to Yassin and Mohamed study because the control chart of Yassin and Mohamed has a lot of subgroups out of control and also shows that the value of ARL_1 is greater than our study result, so in this case, the preference is given to the smallest value. In future work, we can use another control chart with a new approach and use another methodology to treat multicollinearity problems, or we can also use missing values and draw control charts. A lot of ideas can be used in this field.

Disclaimer (Artificial intelligence)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Competing Interests

Authors have declared that no competing interests exist.

References

- Amiria, A., Yeh, B. A., & Asgari, A. (2016). Monitoring two-stage processes with binomial data using generalized linear model-based control charts. *Quality Technology & Quantitative Management*, 13(3), 241–262.
- Bilen, C., Chen, X., Khan, A., & Yadav, P. O. (2007). Multiple regression control chart integrated with principal component analysis. In G. Bayraksan (Ed.), *Industrial Engineering Research Conference*.
- Dunn, K. B., & Smyth, K. G. (2018). *Generalized linear models with examples in R* (pp. 112–130). Springer Science+Business Media, LLC.
- Filho, M. D., & Sant'Anna, O. M. A. (2016). Principal component regression-based control charts for monitoring count data. *The International Journal of Advanced Manufacturing Technology*, 85(5–8), 1565–1574. Available: <https://doi.org/10.1007/s00170-015-8054-6>
- Haridy, S., & Benneyan, J. C. (2024). Shewhart-EWMA chart for monitoring binomial data subject to shifts of random amounts. *Computers & Industrial Engineering*, 193, 467–485. Available: <https://doi.org/10.1016/j.cie.2024.110252>
- Mongkoltawat, P., Areepong, Y., & Sukparungsee, S. (2024). Average run length computations of autoregressive and moving average processes using the extended EWMA procedure. *WSEAS Transactions on Mathematics*, 23, 371–384.

- Nancy, M., Joshi, H., & Dhandra, V. B. (2023). Regression control charts: A survey. *Journal of Pharmaceutical Negative Results*, 14(3), 1079–1086.
- Pardo, A. J., Pardo, L., & Pardo, C. D. M. (2007). A simulation study of a nested sequence of binomial regression models. *Statistics*, 41(3), 253–267.
- Phanyaem, S. (2024). Precise average run length of an exponentially weighted moving average control chart for time series model. *Thailand Statistician*, 22(4), 909–925.
- Rashid, A. N., Mokhtar, F. S., Wan Hassan, S. W., & Che Hussin, E. W. (2013). Regression residual control chart for monitoring academic performance of students in higher learning institution. In *International Conference on Computing, Mathematics and Statistics*.
- Soares, G., Gomes, S., & Ludermir, B. T. (2006). Feature selection for neural networks through binomial regression. In *Neural Information Processing: 13th International Conference, Hong Kong, China, October 3-6* (pp. 737–745). Springer-Verlag Berlin Heidelberg.
- Wilson, J. R., Lorenz, K. A., & Selby, L. P. (2024). Introduction to binary logistic regression. In *Modeling binary correlated responses: Using SAS, SPSS, R and STATA* (pp. 3–18).
- Yassin, S. M., & Mohamed, S. M. (2022). Performance comparison of residual control charts for count data based on ridge regression. *Information Sciences Letters*, 11(1), 2301–2326.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/124816>